



Pediatric Dosing for Ibuprofen (*Motrin*)

The following sheet provides dosing information for a variety of ibuprofen products for infants and children. It is a general guideline and unless otherwise directed by your provider, you should always refer to the directions provided with the product, as they may change.

Dose: Every 6-8 hours as needed. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
If possible, use weight to dose; otherwise, use age.








Infants' Ibuprofen: Concentrated Drops (Motrin)

Active ingredient: Ibuprofen (NSAID*) 50 mg (in each 1.25 mL)

Weight	<i>If child is under 12 lbs. or under 6 months of age, DO NOT administer this medication.</i>	12-17 lbs.	18-23 lbs.
Age		6-11 mos.	12-23 mos.
Dose		 1.25 mL	 1.875 mL






Children's Ibuprofen: Oral Suspension (Motrin)

Active ingredient: Ibuprofen (NSAID*) 100 mg (in each 5 mL)

Weight	<i>If child is under 12 lbs. or under 6 months of age, DO NOT administer this medication.</i>	12-17 lbs.	18-23 lbs.	24-35 lbs.	36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Age		6-11 mos.	12-23 mos.	2-3 years	4-5 years	6-8 years	9-10 years	11 years
Dose		 2.5 mL	 4 mL	 5 mL	 7.5 mL	 10 mL	 12.5 mL	 15 mL





Children's Ibuprofen: Chewables 100mg/tablet (Motrin)

Active ingredient: Ibuprofen (NSAID*) 100 mg (in each chewable tablet)

Weight		24-35 lbs.	36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Age	<i>If child is under 24 lbs. or under 2 years of age, DO NOT administer chewable tablets.</i>	2-3 years	4-5 years	6-8 years	9-10 years	11 years
Dose		1 tablet 	1.5 tablets 	2 tablets 	2.5 tablets 	3 tablets 

Ibuprofen: 200 mg tablets (Motrin)

Active ingredient: Ibuprofen (NSAID*) 200 mg (in each tablet)

Weight		36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Age	<i>If child is under 36 lbs. or under 4 years of age, DO NOT administer tablets.</i>	4-5 years	6-8 years	9-10 years	11 years
Dose		0.5 tablet 	1 tablet 	1 tablet 	1.5 tablets 

* Nonsteroidal anti-inflammatory drug.

If you are unsure about these directions, please contact your Oregon Medical Group clinic and ask to speak to a nurse. We are always happy to help our patients.

Pediatric Dosing for Acetaminophen (TYLENOL®)

The following sheet provides dosing information for a variety of acetaminophen products for infants and children. It is a general guideline and unless otherwise directed by your provider, you should always refer to the directions provided with the product, as they may change.

Dose: Every 4 hours as needed. **DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.** If possible, use weight to dose; otherwise, use age. **DO NOT** use with any other product containing acetaminophen.

Infants' and Children's Acetaminophen: Oral Suspension (Tylenol)

Active ingredient: Acetaminophen 160 mg (in each 5 mL)

Weight	6-11 lbs.	12-17 lbs.	18-23 lbs.	24-35 lbs.	36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Age	0-3 mos.	4-11 mos.	12-23 mos.	2-3 years	4-5 years	6-8 years	9-10 years	11 years
Dose	1.25 mL	2.5 mL	3.75 mL	5 mL	7.5 mL	10 mL	12.5 mL	15 mL

Children's Acetaminophen: Chewables 160 mg/tablet (Tylenol)

Active ingredient: Acetaminophen 160 mg (in each chewable tablet)

Weight		24-35 lbs.	36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Age	If child is under 24 lbs. or under 2 years of age, DO NOT administer chewable tablets.	2-3 years	4-5 years	6-8 years	9-10 years	11 years
Dose		1 tablet	1.5 tablets	2 tablets	2.5 tablets	3 tablets

Acetaminophen: 325 mg tablets (Tylenol)

Active ingredient: Acetaminophen 325 mg (in each tablet)

Weight		24-35 lbs.	36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Age	If child is under 24 lbs. or under 2 years of age, DO NOT administer tablets.	2-3 years	4-5 years	6-8 years	9-10 years	11 years
Dose		Use liquid or chewable for more accurate dosing.		1 tablet	1 tablet	1.5 tablets

If you are unsure about these directions, please contact your Oregon Medical Group clinic and ask to speak to a nurse. We are always happy to help our patients.